

Lyapunov vs. Geometrical Stability Analysis of the Kepler and the Restricted Three Body Problem

A. Yahalom¹, J. Levitan^{2,3*}, M. Lewkowicz² and L. Horwitz^{2,4,5}

¹*Department of Electrical and Electronic Engineering,
Ariel University Center of Samaria, Ariel 40700, Israel*

²*Applied Physics Department, Ariel University Center of Samaria, Ariel 40700, Israel*

³*Department of Physics, Technical University of Denmark, Lyngby 2800, Denmark*

⁴*School of Physics, Tel Aviv University, Tel Aviv 69978, Israel*

⁵*Department of Physics, Bar-Ilan University, Ramat Gan 52900, Israel*

(Dated: July 30, 2009)

In this letter we show that although the application of standard Lyapunov analysis predicts that completely integrable Kepler motion is unstable, the geometrical analysis of Horwitz et al [5] predicts the observed stability. This seems to us to provide evidence for both the incompleteness of the standard Lyapunov analysis and the strength of the geometrical analysis. Moreover, we apply this approach to the three body problem in which the third body is restricted to move on a circle of large radius which induces an adiabatic time dependent potential on the second body. This causes the second body to move in a very interesting and intricate but periodic trajectory; however, the standard Lyapunov analysis, as well as methods based on the parametric variation of curvature associated with the Jacobi metric, incorrectly predict chaotic behavior. The geometric approach predicts the correct stable motion in this case as well.

PACS numbers: 45.20.Jj, 47.10.Df, 05.45.-a, 05.45.Gg

Several relatively recent papers have used a geometric approach to describe Hamiltonian chaos, implementing tools from Riemannian geometry [1-3]. The natural motions of Hamiltonian systems are viewed as geodesics on a Riemannian space associated with a metric g (often associated with either the Jacobi or the Eisenhart metric). Stable motions are thus defined by the curvature properties of the manifold. Casetti et al [2] derived from the Jacobi-Levi-Civita (JLC) equation an effective stability equation which formally describes a stochastic oscillator. They conjectured that some 'average' global geometric property should provide information about the degree of chaos and applied the geometric method to the Fermi-Pasta-Ulam beta-model and to a chain of coupled oscillators. Safaai and Saadat [4] have used the geometric method developed in [1-3] in an attempt to predict chaos in the restricted three body problem. They conclude that the fluctuations of the curvature of the manifold along the geodesics yield parametric instability of the trajectories and cause chaos. They also calculated a positive Lyapunov exponent. It has recently been shown [5] that there is a possibility to characterize instability in Hamiltonian systems by a geometrical approach which takes its point of origin in the curvature associated with a conformal Riemann metric tensor which is applicable to a large class of potential models. This approach appears to be more sensitive than computing Lyapunov exponents or the use of the Jacobi metric. This enables one to associate instability with a negative dynamical curvature appropriate for the geodesic motion different from that defined by the Jacobi metric, for which the line element is proportional to the time τ and not, as for the Jacobi metric, the action. This establishes a natural connection between chaotic Hamiltonian flows and Anosov flows [6].

The geometrical method that we shall use [5] studies the stability of a Hamiltonian system of the form (we use the summation convention)

$$H = \frac{p^2}{2M} + V(x) \quad (1)$$

where V is a function of space variables alone, by introducing a second Hamiltonian of the type considered by Gutzwiller [7, 8]

$$H = \frac{1}{2M} g_{ij} p^i p^j \quad (2)$$

where g_{ij} is a function of the coordinates alone.

Hamilton's equations applied to Eq. (2) result in

$$\ddot{x} = -\Gamma_l^{mn} \dot{x}_m \dot{x}_n ; \quad (3)$$

where

$$\Gamma_l^{mn} = -\frac{1}{2} g_{lk} \left\{ \frac{\partial g^{km}}{\partial x_n} + \frac{\partial g^{kn}}{\partial x_m} - \frac{\partial g^{mn}}{\partial x_k} \right\} \quad (4)$$

and g^{ij} is the inverse of g_{ij} .

Horwitz, et al [5] take g_{ij} to be of conformal form

$$g_{ij} = \frac{E}{E - V(x)} \delta_{ij} \quad (5)$$

on the hypersurface defined by $H = E = \text{const}$, resulting in a formal equivalence between the two Hamiltonians on the same energy shell E .

The velocity field defined by

$$u^j = \dot{x}^j = g^{ij} \dot{x}_i = \frac{p^j}{M} \quad (6)$$

satisfies the geodesic equation

$$\dot{u}^l = -M_{mn}^l u^m u^n \quad (7)$$

where

$$M_{mn}^l \equiv \frac{1}{2} g^{lk} \frac{\partial g_{mn}}{\partial x_k} \quad (8)$$

It is shown in [5] that in the coordinates for which g_{ij} has the form (5), (8) coincides with the Hamilton equations obtained from the Hamiltonian form (1), and thus provides a geometrical embedding of this

Hamiltonian motion. The geodesic deviation ξ^l between two such orbits $x'(t)$ and $x(t)$ satisfies

$$\frac{D_M^2}{D_M t^2} \xi^l = R_{Mqmr}^l \dot{x}^q \dot{x}^n \xi^m \quad (9)$$

where the covariant derivative of ξ^l is defined by

$$\xi^l{}_{;n} = \frac{\partial \xi^l}{\partial x^n} + M_{nm}^l \xi^m \quad (10)$$

$\frac{D_M}{D_M t}$ is the corresponding covariant derivative along the geodesic curve, and

$$R_{Mqmn}^l = \frac{\partial M_{qm}^l}{\partial x^n} - \frac{\partial M_{qn}^l}{\partial x^m} + M_{qm}^k M_{nk}^l - M_{qn}^k M_{mk}^l \quad (11)$$

With the conformal metric in noncovariant form (5), the geodesic deviation equation (9) becomes

$$\frac{D_M^2}{D_M t^2} \xi^l = -\mathcal{V} \xi^l \quad (12)$$

where the matrix $\mathcal{V}$ is given by

$$\mathcal{V}_{li} = \frac{3}{M v^2} \frac{\partial V}{\partial x^l} \frac{\partial V}{\partial x^i} + \frac{1}{M} \frac{\partial^2 V}{\partial x^l \partial x^i} \quad (13)$$

and

$$P^{ij} = \delta^{ij} - \frac{v^i v^j}{v^2} \quad (14)$$

with $v^i = \dot{x}^i$, defining a projection into a direction orthogonal to v^i .

It was argued in [5] that instability should occur if at least one of the eigenvalues of $\mathcal{V}$ is negative. This criterion was found to be effective for many examples [5].

The analysis given above was carried out with a time independent potential function V . We now study the consequences of allowing a weak time dependence in V . In this case E is not precisely conserved, but we shall make an adiabatic approximation in which we consider E to be time independent. The time dependence of V appears in the form of the geodesics generated by the Gutzwiller Hamiltonian (2) as an additional term with partial time derivative of the metric, but in the geometrical imbedding of the Hamiltonian motion, the geodesic evolution of the velocity vector field has the same form (7) as for the time independent potential problem. The formulation of geodesic deviation in terms of the second covariant derivative, however, introduces another time derivative, introducing an additional term in the stability matrix. Following the methods applied above, one finds that the geodesic deviation now obeys the relation

$$\frac{D^2}{Dt^2} \xi^q = R_{Mlrr}^q \dot{x}^l \dot{x}^r \xi^r + \frac{\partial M_{rt}^q}{\partial t} \dot{x}^t \xi^r \quad (15)$$

which is just the result obtained in [5], with the additional term

$$\frac{\partial M_{rt}^q}{\partial t} \dot{x}^t \xi^r = \frac{1}{2} \frac{\partial^2 g_{rt}}{\partial t \partial x_q} \dot{x}^t \xi^r \quad (16)$$

Using the conformal form of g_{ij} with time dependent potential, the right hand side contains

$$\frac{\partial^2 g_{rt}}{\partial t \partial x_q} = \frac{2E}{(E-V(x,t))^3} \frac{\partial V}{\partial t} \frac{\partial V}{\partial x_q} + \frac{E}{(E-V(x,t))^2} \frac{\partial^2 V}{\partial x_q \partial t} \quad (17)$$

Near the boundary of the physical region, these terms can become large, but they are generally small away from these boundaries since they involve the time derivative of V , in the same approximation in which E is constant (the consistency of this approximation will be discussed elsewhere). In our computations here these conditions hold, and we therefore compute stability using the same formulas as for the time independent problem.

We now consider the following Lagrangian of a restricted three body system:

$$L = T - V, \quad T = \frac{1}{2} m_e \left(\frac{d\vec{r}}{dt} \right)^2 \quad (18)$$

$$V = \frac{1}{2} m_j r_j^2 \omega_j^2 - \frac{4\pi^2 m_e}{r} - \frac{4\pi^2 m_j}{r_j} - \frac{4\pi^2 m_e m_j}{r_{12}}$$

In this system three bodies are involved: a large mass M (the "Sun") assumed to be set at the origin (see Fig.1), a mass m_e ("Earth") which its trajectory we wish to study, located at $\vec{r}$, and a third mass m_j (a "Jupiter") located at $\vec{r}_j$ circling the sun with angular frequency ω_j such that

$$\vec{r}_j = r_j (\cos(\omega_j t), \sin(\omega_j t)), \quad (19)$$

r_j is assumed to be constant.




FIG. 1: Depiction of the restricted three body problem




FIG. 2: The Kepler orbit ($m_j = 0$)

Finally $\vec{r}_{12}$ is the distance between the "Earth" and "Jupiter". The following parameters are assumed:

$$r_j = 5.203, \quad \omega_j = \frac{2\pi}{11.86}, \quad m_e = 2.96 \cdot 10^{-6},$$

$$m_{j0} = 317.9 \cdot m_e.$$

Two cases are studied: in the first case we ignore Jupiter, $m_j = 0$; (the classical Kepler problem) and in a second case we assume a "large Jupiter" $m_j = 1000m_{j0}$. The trajectories of the earth in those two cases are depicted in Figs.2 and 3 for the initial conditions $\vec{r}(0) = (1,0)$, $\dot{\vec{r}}(0) = (0,1)$.




FIG. 3: The earth orbit for the restricted three body problem with $m_j = 1000m_{j0}$ during 100 time units

We now study the stability of each of the trajectories described in the previous section using the standard Lyapunov analysis and evaluating the eigenvalues of $\mathcal{V}$ in Eq.(13). The eigenvalues of this matrix are supposed to determine the stability of the trajectory. A negative eigenvalue indicates an unstable trajectory while a positive eigenvalue indicates stability. It can be easily seen that the Lyapunov exponent depends on the eigenvalues of the stability matrix (denoted by $\lambda_{Lyapunov}$ in the figures)

$$\mathcal{V}_{kl} = \frac{1}{m_e} \frac{\partial^2 V}{\partial x_k \partial x_l} \quad k, l \in \{1, 2\} \quad x_1 \equiv x, \quad x_2 \equiv y; \quad (20)$$

the square of the Lyapunov eigenvalues are given by the negative of the eigenvalues of this matrix.

In the geometrical approach we write the matrix $\mathcal{V}$; Eq.(13), as:

$$\mathcal{V}_{kl} = \frac{1}{m_e} \left[\frac{3}{2(E-V)} \frac{\partial V}{\partial x_k} \frac{\partial V}{\partial x_l} + \frac{\partial^2 V}{\partial x_k \partial x_l} \right], \quad (21)$$

where E is now assumed to be adiabatically constant.

For the Kepler case, one eigenvalue is always positive; the second eigenvalue is depicted in Fig.4. Clearly it becomes negative during certain short time intervals. We found that this occurs at the perihelia of the orbits. A local Lyapunov instability is not a cause for concern since the orbit may be globally stable. However, Fig.4 reveals that the Lyapunov eigenvalue is indeed negative along the entire orbit although its negative value is less pronounced between perihelia than at the perihelia. Thus according to Lyapunov analysis the Kepler problem is unstable and could become chaotic. This is clearly a false result since the Kepler problem is integrable.

Yet the "geometric eigenvalue", depicted in Fig.5, is found to be positive in between perihelia, hence the trajectory is predicted to be globally stable. The trajectory only spends a short time at the unstable perihelia and thus instability does not have enough time to take place. This is a clear indication that the geometrical analysis is preferable.

The trajectory of the restricted three body problem is shown in Fig.3. This trajectory is quite complicated, however, it appears to be periodic rather than chaotic. The Lyapunov eigenvalues and the geometric eigenvalues are depicted in Fig.6 and Fig.7.

The eigenvalues in the restricted three body problem have a different behaviour than those of the Kepler problem. Nevertheless, the same conclusions are obtained: Lyapunov analysis predicts instability and possible chaos, geometrical analysis predicts stability. The application by Safaai et al [4] of the methods of Cassetti and Pettini also appear to predict chaotic behaviour, which is not in agreement with our simulations.




FIG. 4: The eigenvalues of the Lyapunov stability matrix (Eq.(20)) for Kepler problem, $m_j = 0$. Upper left: during 1 time unit; upper right: passing a perihelion; middle: between perihelia.




FIG. 5: The eigenvalues of the "Geometry" stability matrix (Eq.(21)) for Kepler problem, $m_j = 0$. Upper left: during 1 time unit; upper right: passing a perihelion; middle: between perihelia.




FIG. 6: The eigenvalues of the Lyapunov stability matrix (Eq.(20)) for the restricted three body problem with $m_j = 1000m_{j0}$. Upper left: during 1 time unit; upper right: passing a perihelion; middle: between perihelia.




FIG. 7: The eigenvalues of the "Geometry" stability matrix (Eq.(21)) for the restricted three body problem with $m_j = 1000m_{j0}$. Upper left: during 1 time unit; upper right: passing a perihelion; middle: between perihelia.

*Electronic address: levitan@ariel.ac.il

- [1] M. Pettini, Phys. Rev. E 47, 828 (1993).
- [2] L. Casetti and M. Pettini, Phys. Rev. E 48, 4320 (1993).
- [3] M. Cerutti-Sola and M. Pettini, Phys. Rev. E 51, 53 (1994).
- [4] H. Safaai, M. Hasan and G. Saadat, *Understanding Complex Systems* (Springer Berlin, 2006).
- [5] L. Horwitz, Y. Ben Zion, M. Lewkowicz, M. Schiffer and J. Levitan, Phys. Rev. Lett, 98, 234301 (2007). See also Y. Ben Zion and L. Horwitz, Phys. Rev. E 76, 046220 (2007); Y. Ben Zion and L. Horwitz, Phys. Rev. E 78, 036206 (2008).
- [6] D. Anosov, Geodesic flows on closed Riemannian manifolds with negative curvature, Proc Steklov Instit. Math. 90, Amer. Math. Soc. Providence, Rhode Island, 1969.

- [7] C.G.J. Jacobi, *Vorlesungen über Dynamik*, Verlag G. Reiner, Berlin (1844); J.S. Hadamard, J. Math. Pures et Appl. 4, 27 (1898).
- [8] M.C. Gutzwiller, *Chaos in Classical and Quantum Mechanics*, Springer-Verlag, New York (1990). See also W.D. Curtis and F.R. Miller, *Differentiable Manifolds and Theoretical Physics*, Academic Press, New York (1985), J. Moser and E.J. Zehnder, *Notes on Dynamical Systems*, Amer. Math. Soc., Providence (2005), and L.P. Eisenhardt, *A Treatise on the Differential Geometry of Curves and Surfaces*, Ginn, Boston (1909) [Dover, N.Y. (2004)].